





To: Indira IVF Hospital Pvt Ltd-Dehradun

Ashirwad Towers, Above Nexa Showroom Ram

Vihar,

Ballupur Road, Chowk, Chakarata Rd, Ballupur,

Uttara Khand

Dehradun - 248001

Contact:

Report Of: Mrs. PAPLI SISODIA

Pt. Contact: 8171308844



Sample ID	2100119433			
Patient ID	1002187732			
Received on	10/12/2021 15:54			
Registered on	11/12/2021 14:56			
Reported on	15/12/2021 22:01			
Referred by	DR.REEMA SIRCAR			
Sonography by	DR.RAGHAV BHATIA			

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. PAPLI SISODIA Patient DOB: 03/08/1977

Ethnicity: Asian City: DEHRADUN Hospital ID: FDUKDC20

Sample Type: Serum Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method: Time-resolved Fluroimmunoassay

EVIC Screen* is an evidence based prenatal screening program curated by Lilac Insights in accordance with the international guidelines for prenatal screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 5 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies
- External aExternal audit of the prenatal screening program by United Kingdom National External Quality Assessment Service (UKNEQAS) scheme and Randox International Quality Assessment Scheme (RIQAS)

RISK ASSESSMENT				
T21 (Down syndrome)	1: 29605	Low Risk	LOW	INTERMEDIATE
T18 (Edwards' syndrome)	1: 100000	Low Risk	LOW	HIGH
T13 (Patau syndrome)	1: 32058	Low Risk	LOW	HIGH

MULTIPLE OF MEDIAN (MoM)							
Free ß-hCG	0.64						
AFP	2.29						
PAPP-A	0.85						
PLGF	1.09						

INTERPRETATION

The First Trimester - Quadruple Screening for the given sample is found **SCREEN NEGATIVE**.

SUGGESTIONS AND OTHER FINDINGS


In view of increased Alpha-fetoprotein (AFP), detailed anomaly scan to assess for fetal abnormalities especially that of the spine, anterior abdominal wall, and kidneys.







Verified by
Mr. Pradip Kadam
Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path), Consultant Pathologist









Patient name: Mrs. PAPLI SISODIA Sample ID: 2100119433

			PREGNANC	Y DETAILS				
No. of fetuse:	s :1		EDD	: 24/06/2022	Age at Term	: 30.4	Years	
GA is Based on : Ass. rep.		LMP Date	: 18/09/2021	LMP Certainty: Regular		lar		
Smoking: No	Smoking: None Parity:		Height	:	Weight : 60.60 Kg		0 Kg	
FHR :								
Pr	revious pregnancy histo	ory	Pre-ec	Pre-eclampsia history Other findi			dings	
Down sy	yndrome Edwards's	syndrome	PE in pre	Insuli	Insulin dependent diabetes			
Patau sy	ndrome NTD synd	rome	Pat. mot	Chro	Chronic hypertension			
Assisted Rep	roduction : Donor egg Ti	ransfer Date :	04/10/2021 E	Extraction Date: 30/0	1/2021 D ono	r DOB : 15	/06/1991	
Note! Age at 1	term is calculated from the	Donor DOB						
EDD: Estimate	ed Due Date GA: Gestation Age	LMP: Last Mer	nstrual Period FHI of Bir	•	Neural Tube Defe	ct PE: Pre-e	clampsia DOB: Date	
			SPECIMEN					
			SPECIMEN	DE IAILS				
Sample ID	:2100119433	CRL :5	0.5 mm	Test Name	Conc.	Unit	Corr. Mom	
Collection Da	ate :08/12/2021	CRL2 :		Free-ß-hCG	31.02	ng/mL	0.64	
Scan Date	:08/12/2021	BPD :		AFP	20.07	U/mL	2.29	
GA at Coll Da	ate: 11 Weeks 5 Days	BPD2 :		PAPP-A	2061.00	mU/L	0.85	
GA at Scan D	ate: 11 Weeks 5 Days	HC :		PLGF	42.78	pg/mL	1.09	
Received on	:10/12/2021	HC2 :						
GA: Gestation	Age CRL: Crown Rump Length	BPD: Bi-parie	tal Diameter HC: I	Head Circumference free	-ß-hCG: free-Beta l	Human Chor	ionic Gonadotropin	
	NT: I	Nuchal Transluc	cency PAPP-A: Pre	gnancy-associated Plasma	Protein-A			
RISKS								
Disorder: Down Syndrome				Res	Result: Low Risk			
Final risk:	1:29605	Age risk:	1:926					
Cutoff	1:250	Risk type	Risk At Term					
Disorder: Edwards' Syndrome				Res	sult:	Low Risk		
Final risk:	1:100000	Age risk:	1:8333					
Cutoff	1:100	Risk type	Risk At Term					
Disorder: Pat	Disorder: Patau Syndrome Result: Low Risk							
Final risk:	1:32058	Age risk:	1:25021					
Cutoff	1:100	Risk type	Risk At Term					











Patient name: Mrs. PAPLI SISODIA Sample ID: 2100119433

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Intermediate

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- Quality of the Down's Syndrome & ONTD screening program (Biochemical values, MoMs and Risk assessments) monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- $\bullet \quad \text{Lilac Insights does not bear responsibility for the Ultra sound measurements.}\\$
- This is a risk estimation test and not a diagnostic test. An increased risk result does not mean that the fetus is affected and a low risk result does not mean that the fetus is unaffected. Reported risks should be correlated and adjusted according to the absence/presence of sonographic markers observed in the anomaly/malformation scan.
- The above risk has been calculated based on Biochemistry values alone.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and further tests must be performed before a firm diagnosis can be made. A low risk result does not exclude the possibility of Down's Syndrome or other abnormalities, as the risk assessment does not detect all affected pregnancies.





END OF REPORT

