



To: Indira IVF Hospital Pvt Ltd-Bareilly

Plot No-1st & 2nd Floor

Nijhawan Tower, Station Road, Opp -ADM(E)

Compound, Civil Lines

Bareilly - 243001 Contact: 7042094494

Report Of: Mrs. MANJU GAUTAM

Pt. Contact: 9456869966



Sample ID 2200004554

Patient ID 1002199475

Received on 10/01/2022 14:15

Registered on 12/01/2022 12:03

Reported on 12/01/2022 20:00

Referred by DR.SHIVI SAXENA

Sonography by DR.SUMIT TANDAN

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. MANJU GAUTAM Patient DOB: 20/03/1980

Ethnicity: Asian City: BAREILLY Hospital ID: GBUPFD15

Sample Type: Serum

Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method: Time-resolved Fluroimmunoassay

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 5 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies
- External audit of the prenatal screening program by United Kingdom National External Quality Assessment Service (UKNEQAS) scheme and Randox International Quality Assessment Scheme (RIQAS)

RISK ASSESSMENT Low Risk T21 (Down syndrome) 1:5013 Low Risk 1:4997 T18 (Edwards' syndrome) 1:100000 Low Risk 1:100000 Low Risk T13 (Patau syndrome) 1:100000 1:100000 Low Risk Low Risk MULTIPLE OF MEDIAN (MoM Free ß-hCG 3.51 PAPP-A 1.28

INTERPRETATION

The First Trimester Screening for the given sample is found **SCREEN NEGATIVE**.





Verified by
Mr. Pradip Kadam

Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path). Consultant Pathologist





Patient name: Mrs. MANJU GAUTAM Sample ID: 2200004554

PREGNANCY DETAILS							
No. of fetuses	:2DCDA	EDI)	: 18/07/2022	Age at Term	n :31.4	Years
GA is Based on	: Ass. rep.	LMI	P Date	: 14/10/2021	LMP Certai	inty : Regu	ılar
Smoking: None	Parity :	Hei	ght	:	Weight	: 59.5	0 Kg
FHR:							
Previous pregnancy history Pre-eclampsia history Other findings							ndings
Down syndrom	ne Edwards' syn	drome	PE in previous pregnancy		Insulin dependent diabetes		
Patau syndrom	e NTD syndron	ne	Pat. motl	her had PE	Chronic hypertension		
Assisted Reproduction : Donor egg Transfer Date : 28/10/2021 Extraction Date : 23/10/2021 Donor DOB : 04/02/1991 Note! Age at term is calculated from the Donor DOB							
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date							
of Birth							
SPECIMEN DETAILS							
Sample ID :	: 2200004554	CRL : 60 mm	1	Test Name	Conc.	Unit	Corr. Mom
Collection Date :	:07/01/2022	CRL2 :61 mm	า	Free-ß-hCG	291.20	ng/mL	3.51
Scan Date :	:07/01/2022	BPD :		NT	1.2	mm	0.78
GA at Coll Date : 12 Weeks 4 Days BPD2 :		BPD2 :	NT2		1.25	mm	0.80
GA at Scan Date :	: 12 Weeks 4 Days	HC :		PAPP-A	9120.00	mU/L	1.28
Received on :	: 10/01/2022	HC2 :					
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A							
RISKS							
Disorder: Down Syn	drome			Result:		Result:	
Twin 1 Twin 2		n 2			Twin 1		Twin 2
Final risk: 1:501		1:4997	Age risk:	1:669	Low Risk	Low	
Cutoff: 1:250	Cutoff:	1:250	Risk type	: Risk At Term			
Disorder: Edwards' Syndrome Result: Result:							Result:
Twin 1 Twin 2					Twin 1		Twin 2
Final risk: 1:100		1:100000	Age risk:	1:3614	Low Risk	Low	
Cutoff: 1:100		1:100	Risk type		LOW KISK	Low	KISK
··							
Disorder: Patau Syndrome Twin 1 Twin 2					Result:		Result:
Twin 1 Final risk: 1:100		1:100000	Age risk:	1:10851	Twin 1		Twin 2
Cutoff: 1:100		1:100	Risk type		Low Risk	Low	KISK







Patient name: Mrs. MANJU GAUTAM Sample ID: 2200004554

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk **Intermediate Risk result:** An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- $\bullet \quad \text{Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.}\\$
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for the NT & CRL measurements. We strongly recommend that NT/ CRL measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.

END OF REPORT



