





To: Midtown Diagnostics

Navneelam Building, 2nd Floor,

Above CCD, First Right After Potdar Hospital, Worli

Maharashtra

Mumbai - 400018

Contact: 022-24976302

Report Of: Mrs. ARWA KAPOOR

Pt. Contact:

Sample ID 2200083519

Patient ID

Received on 30/07/2022 10:07

1002254216

Registered on 30/07/2022 15:26

Reported on 04/08/2022 10:32

Referred by DR.A PANDOLE

Sonography by DR.ROMA LALA

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. ARWA KAPOOR Patient DOB: 16/09/1984

Ethnicity: Asian City: MUMBAI Hospital ID:

Sample Type:Serum

Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method:Time-resolved Fluroimmunoassay

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 5 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies
- External audit of the prenatal screening program by United Kingdom National External Quality Assessment Service (UKNEQAS) scheme and Randox International Quality Assessment Scheme (RIQAS)

RI	SK ASSESSME	NT		
T21 (Down syndrome)	1: 100000	Low Risk	LOW	INTERMEDIATE HIGH
T18 (Edwards' syndrome)	1:100000	Low Risk	LOW	HIGH
T13 (Patau syndrome)	1: 100000	Low Risk	LOW	HIGH
Pre-eclampsia before 34 wee	ks 1:4539	Low Risk	LOW	HIGH
Pre-eclampsia before 37 wee	ks 1:446	Low Risk	LOW	HIGH

MULTIPLE OF MEDIAN (MoM)

Free ß-hCG	0.59	
AFP	1.61	
PAPP-A	1.24	
PLGF	0.97	

INTERPRETATION

The First Trimester Enhanced Screening for the given sample is found **SCREEN NEGATIVE**.

SUGGESTIONS AND OTHER FINDINGS

• Please correlate clinically with USG findings.







Verified by
Mr. Pradip Kadam
Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path), Consultant Pathologist

Page 1 of 3









Patient name: Mrs. ARWA KAPOOR				Sample	eID: 2200083519			
	PREGNANC	Y DETAILS						
No. of fetuses : 1	EDD	: 13/02/2023	Age at Terr	n :38.4	Years			
GA is Based on : Ass. rep.	LMP Date	:	LMP Certainty: Unknown					
Smoking: None Parity: Nulliparous	Height	: 165.1 cm	•		O Kg			
FHR:	•				J			
Previous pregnancy history	Pre-ec	lampsia history		Other fin	dings			
Down syndrome Edwards' syndrome								
Patau syndrome NTD syndrome								
Assisted Reproduction: IVF Transfer Date: 26		Extraction: 38 yrs	Cinc	orne rryper te	.1131011			
			Noural Tuba Daf	ant / DE Dra or	olempsis I DOR: Date			
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date of Birth								
	SPECIMEN	DETAILS						
Sample ID : 2200083519 CRL	: 48.6 mm	Test Name	Conc.	Unit	Corr. Mom			
Collection Date : 29/07/2022 CRL2	:	Free-ß-hCG	29.94	ng/mL	0.59			
Scan Date : 29/07/2022 BPD	:	NB	Present					
GA at Coll Date : 11 Weeks 4 Days BPD2	2 :	AFP	13.78	U/mL	1.61			
GA at Scan Date : 11 Weeks 4 Days HC	:	NT	1.1	mm	0.86			
Received on : 30/07/2022 HC2	:	PAPP-A	2781.00	mU/L	1.24			
		PLGF	38.72	pg/mL	0.97			
		MAP	83.33	mmHg	0.99			
		UTPI	1.04		0.58			
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A								
	RISK	(S						
Disorder: Down Syndrome		Re	sult:	Low Risk				
Final risk: 1:100000 Age risk	: 1:194							
Cutoff 1:250 Risk typ	e Risk At Term							
Disorder: Edwards' Syndrome		Re	sult:	Low Risk				
Final risk: 1:100000 Age risk	: 1:1746		our.	20111131				
Cutoff 1:100 Risk typ								
Disorder: Patau Syndrome		Re	sult:	Low Risk	•			
Final risk: 1:100000 Age risk	: 1:5238		our.	20111131				
Cutoff 1:100 Risk typ								
Disorder: PE < 34 weeks Result: Low Risk								
Final risk: 1: 4539		Ke		_0 14.514				
Cutoff 1: 100 Risk typ	e Risk at Term							
Disorder: PE < 37 week		Re	sult:	Low Risk	•			
Final risk: 1: 446		KC.		20 11 1131				
Cutoff 1: 100 Risk typ	e Risk at Term				Page 2 of 3			



Cutoff



1:100



Risk at Term

Risk type













Patient name: Mrs. ARWA KAPOOR Sample ID: 2200083519

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk **Intermediate Risk result:** An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- $\bullet \quad \text{Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.}$
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- PE risk stratification is done using a cut-off of 1:100 as per ASPRE study.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.

END OF REPORT Page 3 of 3





