





To: Beams Multispeciality Hospital-Khar West

Chrysalizz Building, CD Marg, Plot no. 67,

Between 18th and 19th Road, Khar Danda Road,

Khar West

Maharashtra

Mumbai - 400052

Contact: 022-68188888

Report Of: Mrs. SHRADDHA DANDEKAR

Pt. Contact: 8169147641

Sample ID 2200087138

Patient ID 1002255702

Received on 02/08/2022 21:07

Registered on 03/08/2022 18:04

Reported on 04/08/2022 10:57

Referred by DR.MANJIRI KAWDE

Sonography by DR.SANJEEV MANI

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. SH	RADDHA DANDEKAR	Patient DOB: 12/04/1993	
Ethnicity: Asian	City: MUMBAI	Hospital ID:	

Sample Type:Serum


Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method: Electrochemiluminescence

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 5 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies
- External audit of the prenatal screening program by United Kingdom National External Quality Assessment Service (UKNEQAS) scheme and Randox International Quality Assessment Scheme (RIQAS)

RI	SK ASSESSME	NT		
T21 (Down syndrome)	1:5100	Low Risk	LOW	INTERMEDIATE HIGH
T18 (Edwards' syndrome)	1:100000	Low Risk	LOW	HIGH
T13 (Patau syndrome)	1:80000	Low Risk	LOW	HIGH



INTERPRETATION


The First Trimester Screening for the given sample is found SCREEN NEGATIVE.

SUGGESTIONS AND OTHER FINDINGS

In view of the raised serum free β hCG, fetal growth scan is suggested at 28 - 30 weeks in addition to their routine antenatal care.







Verified by **Mr. Pradip Kadam** Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path), Consultant Pathologist











Patient name: Mrs. SHRADDHA DANDEKAR Sample ID: 2200087138

				PRI	EGNANC	/ DETAILS				
No. of fetuse	lo. of fetuses : 1		EDI)	: 18/02/2023		Age at Term : 29.8 Years		Years	
GA is Based o	GA is Based on : CRL 46.1mm at 01/08/2022		LMI	P Date	: 14/05/2022		LMP Certainty : Regular		lar	
Smoking: No	Smoking: None Parity:		Hei	ght	:		Weight : 86.60 Kg			
FHR :										
Previous pregnancy history				Pre-eclampsia history Other 1			Other fin	dings		
Down syndrome Edwards' syndrome			_, <u></u>	PE in previous pregnancy			Insulin dependent diabetes			
Patau syndrome NTD syndrome				Pat. mother had PE			Chronic hypertension			
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date									clampsia DOB: Date	
of Birth									, ,	
				SF	PECIMEN	DETAILS				
Sample ID	:	2200087138	CRL	: 46.1 ı	mm	Test Name		Conc.	Unit	Corr. Mom
Collection D	ate :	01/08/2022	CRL2	:		Free-ß-hCG		107.40	ng/mL	3.20
Scan Date	:	01/08/2022	BPD	:		NB		Present		
GA at Coll Da	ate :	11 Weeks 2 Days	BPD2	:		NT		1.2	mm	1.08
GA at Scan D	Date :	11 Weeks 2 Days	HC	:		PAPP-A		946.00	mIU/L	0.82
Received on	:	02/08/2022	HC2	:						
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin									ionic Gonadotropin	
NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A										
					RISK	S				
Disorder: Down Syndrome						Resu	lt:	Low Risl		
Final risk:	1:5100		Age risk:	1:1	1000					
Cutoff	1:250		Risk type	Ris	sk At Term					
Disorder: Edwards' Syndrome						Resu	lt:	Low Risl	(
Final risk:	1:1000	00	Age risk:	1:7	7500					
Cutoff	1:100		Risk type	Ris	sk At Term					
Disorder: Patau Syndrome Result:					lt:	Low Risl				
Final risk:	1:8000	C	Age risk:	1:1	11000					_
Cutoff	1:100		Risk type	Ris	sk At Term					



Cutoff





Risk type











Patient name: Mrs. SHRADDHA DANDEKAR Sample ID: 2200087138

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

i i

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- $\bullet \quad \text{Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.}\\$
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.

END OF REPORT

Page 3 of 3





