



To: Apollo Multispeciality Hospital-Kolkata

58, Canal Circular Road, Kadapara Phool Bagan Kolkata - 700054 Contact: 7903876206

Report Of: Mrs. NEELAM SHARMA

Pt. Contact:



 Sample ID
 2300027438

 Patient ID
 10022139044

 Received on
 16/02/2023 11:38

 Registered on
 20/02/2023 17:54

 Reported on
 21/02/2023 16:46

 Referred by
 DR.R. BANERJEE

 Sonography by
 DR.S.R PAL

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. NEELAM SHARMA Patient DOB: 25/06/1991

Ethnicity: Asian City: KOLKATA Hospital ID: AMHLOPP5584366

Sample Type:DBS

Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method:Time-resolved Fluroimmunoassay

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 5 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies
- External audit of the prenatal screening program by United Kingdom National External Quality Assessment Service (UKNEQAS) scheme and Randox International Quality Assessment Scheme (RIQAS)

RI				
T21 (Down syndrome)	1:5792	Low Risk	LOW	INTERMEDIATE HIGH
T18 (Edwards' syndrome)	1: 12107	Low Risk	LOW	HIGH
T13 (Patau syndrome)	1: 15254	Low Risk	LOW	HIGH

MULTIPLE OF MEDIAN (MoM)									
Free ß-hCG	0.69								
PAPP-A	0.30								

INTERPRETATION

The First Trimester Screening for the given sample is found **SCREEN NEGATIVE**.

SUGGESTIONS AND OTHER FINDINGS

In view of PAPP-A MoMs observed in the mother, focused serial survillance for assessment of fetal growth and possiblity of other rare chromosomal/gene defect. Development of high blood pressure related problems in the mother can be considered.







Incharae Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path), Consultant Pathologist

of 3





Patient name: Mrs. NEELAM SHARMA Sample ID: 2300027438

PREGNANCY DETAILS											
No. of fetuse	:s	:1		EDD	: 25/08/2023	Age at Terr	n :32.2	Years			
GA is Based on : CRL 63.1mm at 15/02/2023		LMP Date	: 17/11/2022	LMP Certainty : Regular		lar					
Smoking: None Parity:		Height	:	Weight	Weight : 77.00 Kg						
FHR:											
Pı	reviou	s pregnancy histo	ory	Pre-ecla	ampsia history	Other findings					
Down syndrome Edwards' syndrome			PE in prev	PE in previous pregnancy		Insulin dependent diabetes					
Patau syndrome NTD syndrome			Pat. mother had PE		Chronic hypertension						
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date											
				of Birtl	h						
				SPECIMEN I	DETAILS						
Sample ID		:2300027438	CRL	: 63.1 mm	Test Name	Conc.	Unit	Corr. Mom			
Collection D	ate	: 15/02/2023	CRL2	:	Free-ß-hCG	22.62	ng/mL	0.69			
Scan Date		: 15/02/2023	BPD	:	NB	Present					
GA at Coll Da	ate	: 12 Weeks 5 Days	BPD2	:	NT	1.6	mm	1.00			
GA at Scan D	ate	: 12 Weeks 5 Days	HC	:	PAPP-A	0.26	U/L	0.30			
Received on		: 16/02/2023	HC2	:							
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin											
NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A											
RISKS											
Disorder: Down Syndrome Result: Low Risk							(
Final risk:	1:579	2	Age risk:	1:721							
Cutoff	1:250		Risk type	Risk At Term							
Disorder: Edwards' Syndrome Result: Low Risk											
Final risk:	1:121	07	Age risk:	1:6489							
Cutoff	1:100		Risk type	Risk At Term							
Disorder: Patau Syndrome Result: Low Risk											
Final risk:	1:152	54	Age risk:	1:19480							



1:100

Cutoff



Risk At Term

Risk type







Patient name: Mrs. NEELAM SHARMA Sample ID: 2300027438

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- $\bullet \quad \text{Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.}\\$
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

UK NEQAS
International Quality Expertise
Lab Reg. No. 90968

Page 3 of 3

END OF REPORT