



To: Sampat Clinic-Indore

202 SV Bussiness Park

Madhya Pradesh

Indore - 452001

Contact:

Report Of: Mrs. SWATI AGRAHARI

Pt. Contact: 9589649900



Sample ID 2300102241

Patient ID 1002341101

Received on 15/06/2023 18:41

Registered on 20/06/2023 16:40

Reported on

Referred by

Sonography by

DR.MANJU PATIDAR

DR.SHWETA JAIN

Patient DOB: 02/01/1993

Understand Your Report In Detail Scan OR code

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. SWATI AGRAHARI

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probality of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

UKNEQAS: United Kingdom National External Quality **Assessment Service**

RIQAS: Randox International Quality Assessment Scheme



The Risk Assessment Performed Using CE-Marked Antenatal Risk Evaluation Software Certified by the British Standards Institute (BSI)- ISO 13485:2016

RISK ASSESSMENT T21 (Down syndrome) 1:1146 Low Risk LOW INTERMEDIATE HIGH LOW T18 (Edwards' syndrome) 1:100000 Low Risk HIGH T13 (Patau syndrome) 1:100000 Low Risk LOW HIGH Low Risk Pre-eclampsia before 34 weeks 1:228 HIGH

1ULTIPLE OF MEDIAN (MoM Free ß-hCG 4.09 PAPP-A 0.75

INTERPRETATION

The First Trimester Screening for the given sample is found SCREEN NEGATIVE.


SUGGESTIONS AND OTHER FINDINGS

In view of free bHCG MoMs observed in the mother, kindly consider correlation with fetal growth and well being scan at 28 - 30 weeks.









Verified by Dr. Suresh Bhanushali MD (Path), Consultant Pathologist





Patient name: Mrs. SWATI AGRAHARI Sample ID: 2300102241

Sample Type:Serum Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method: Electrochemiluminescence

Method:Elec	trocne	emiluminescence							
					PREGNANCY	DETAILS			
No. of fetuse	s	:1			EDD	: 20/12/2023	Age at Tern	n :31.0	Years
GA is Based on : CRL 66mm at 13/06		/06/2023		LMP Date : LMP Certainty : Unkno		nown			
Smoking: None Parity: Nulliparous				Height	: 157.4 cm	Weight : 65.00 Kg			
Ethinicity: Asian FHR:									
Previous pregnancy history					Pre-ecla	Other findings			
Down syndrome Edwards' syndrome				PE in previous pregnancy		Insulin dependent diabetes			
Patau syndrome NTD syndrome			•		Pat. mothe	Chronic hypertension			
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date									
of Birth									
SPECIMEN DETAILS									
Sample ID		: 2300102241	CRL	:6	66 mm	Test Name	Conc.	Unit	Corr. Mom
Collection Da	ate	: 14/06/2023	CRL2	:		Free-ß-hCG	115.00	ng/mL	4.09
Scan Date		: 13/06/2023	BPD	:		NB	Present		
GA at Coll Da	ate	: 13 Weeks 0 Days	BPD2	:		NT	1.2	mm	0.81
GA at Scan D	ate	: 12 Weeks 6 Days	HC	:		PAPP-A	3105.00	mIU/L	0.75
Received on		: 15/06/2023	HC2	:		MAP	93.33	mmHg	1.09
						UTPI	1.70		1.11
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin									
NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A									
					RISKS	5			
Disorder: Down Syndrome						Res	Result: Low Risk		
Final risk: 1:1146		Age risk:		1:858					
Cutoff	1:250		Risk type		Risk At Term				
Disorder: Edwards' Syndrome Result: Low Risk									
Final risk: 1:100000		Age risk:		1:7722					
Cutoff	1:100		Risk type		Risk At Term				
Disorder: Patau Syndrome						Res	ult:	Low Risk	(
Final risk: 1:100000		Age risk:		1:23183					
Cutoff	1:100		Risk type		Risk At Term				
Disorder: PE < 34 weeks						Res	ult:	Low Risk	(
Final risk:	1:228	3							
Cutoff	1: 100)	Risk type		Risk at Term				











Patient name: Mrs. SWATI AGRAHARI Sample ID: 2300102241

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Intermediate

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: <u>www.lilacinsights.com/faq-pns</u>

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- PE risk stratification is done using a cut-off of 1:100 as per ASPRE study.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT

Page 3 of 3



