





To: Alpha Women's Clinic-Salem

 $Alpha\,Women's\,Clinic,\,2\hbox{-}a\,/\,131,\,1st\,floor\,Arunagiri$

ENT Hospital

Saradha College Road, Salem.

Tamil Nadu

Salem - 636016

Contact:

Report Of: Mrs. HEMA HARIHARAN

Pt. Contact: 1000000000



Sample ID	2300170383	Understand Your				
Patient ID	1102324100	Report In Detail				
Received on	12/10/2023 13:39					
Registered on	12/10/2023 14:57					
Reported on	-	Scan QR code				
Referred by	Dr. SELVABHARATHY A					
Sonography by	Ramva					

Patient DOB: 07/05/1995

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. HEMA HARIHARAN

EVIC Screen is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probality of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

UKNEQAS: United Kingdom National External Quality Assessment Service

RIQAS: Randox International Quality Assessment Scheme



The Risk Assessment Performed Using
CE-Marked Antenatal Risk Evaluation Software
Certified by the British Standards Institute
(BSI)- ISO 13485:2016

RISK ASSESSMENT LOW T21 (Down syndrome) 1:12000 Low Risk INTERMEDIATE HIGH Free ß-hCG 6.55 T18 (Edwards' syndrome) 1:100000 Low Risk LOW HIGH PAPP-A 2.06 T13 (Patau syndrome) 1:100000 Low Risk LOW HIGH

INTERPRETATION

The First Trimester Screening for the given sample is found **SCREEN NEGATIVE**.

SUGGESTIONS AND OTHER FINDINGS

In view of free bHCG MoMs observed in the mother, kindly consider correlation with fetal growth and well being scan at 28 - 30 weeks.







Verified by **Mr. Pradip Kadam** Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali**MD (Path), Consultant Pathologist











Patient name: Mrs. HEMA HARIHARAN Sample ID: 2300170383

Sample Type: Serum Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method:Electrochemiluminescence											
				PREGNANCY DETAILS							
No. of fetuse	s	:1		EDD	: 17/04/2024	Age at Term	: 28.9	Years			
GA is Based on : CRL 59.3mm at 07/10/2023		LMP Date	: 12/07/2023	LMP Certai	LMP Certainty : Regular						
Smoking: None Parity:		Height	:	Weight	Weight : 73.20 Kg						
Ethinicity: Asian FHR :											
Previous pregnancy history			Pre-ecla	ampsia history		Other findings					
Down syndrome Edwards' syndrome			PE in prev	ious pregnancy	Insul	Insulin dependent diabetes					
Patau syndrome NTD syndrome		•	Pat. mother had PE		Chronic hypertension						
EDD: Estimated Due Date GA: Gestation Age LMP: Last Menstrual Period FHR: Fetal Heart Rate NTD: Neural Tube Defect PE: Pre-eclampsia DOB: Date											
				of Birtl							
SPECIMEN DETAILS											
Sample ID		:2300170383	CRL :	59.3 mm	Test Name	Conc.	Unit	Corr. Mom			
Collection Da	ate	:07/10/2023	CRL2 :		Free-ß-hCG	213.60	ng/mL	6.55			
Scan Date		:07/10/2023	BPD :		NB	Present					
GA at Coll Da	ate	: 12 Weeks 3 Days	BPD2 :		NT	1.3	mm	0.97			
GA at Scan D	ate	: 12 Weeks 3 Days	HC :		PAPP-A	6047.00	mIU/L	2.06			
Received on		: 12/10/2023	HC2 :								
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A											
RISKS											
Disorder: Down Syndrome				Re	Result: Low Risk		(
Final risk:	1:12000 Age risk:		1:1100								
Cutoff	1:250		Risk type	Risk At Term							
Disorder: Edwards' Syndrome				Re	esult:	Low Risk	(
Final risk: 1:100000 Age ris		Age risk:	1:7900								
Cutoff	1:100		Risk type	Risk At Term							
Disorder: Patau Syndrome Result: Low Risk											
Final risk: 1:100000 Age risk:		1:12000									



Cutoff



1:100



Risk At Term

Risk type











Patient name: Mrs. HEMA HARIHARAN Sample ID: 2300170383

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: www.lilacinsights.com/faq-pns

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub-chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT







