





To:	<b>Devi Hospital-Salem</b> No.136/76,New Fairlands,		SampleID	2300170526	Understand Your	
	Tamil Nadu		Patient ID	1102325632	Report In Detail	
	Salem - 636016		Received on	23/10/2023 13:18		
	Contact:		Registered on	23/10/2023 14:45		
	Report Of: Mrs. BHAVANI RAMU		Reported on	24/10/2023 05:30	回新約35%	
	Pt. Contact: 100000000		Referred by	_ , _ ,	Scan QR code	
			,	Dr. M.SRIDEVI		
			Sonography by	Dr. M SRIDEVI		

# EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

## Patient Name: Mrs. BHAVANI RAMU

## Patient DOB: 17/05/1997

EVIC Screen" is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK)

guidelines for First Trimester Screening to determine the probality of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

## UKNEQAS: United Kingdom National External Quality Assessment Service

RIQAS: Randox International Quality Assessment Scheme



The Risk Assessment Performed Using CE-Marked Antenatal Risk Evaluation Software Certified by the British Standards Institute (BSI)- ISO 13485:2016

#### **RISK ASSESSMENT 1ULTIPLE OF** MEDIAN (MoM T21 (Down syndrome) 1:77000 Low Risk LOW INTERMEDIATE HIGH Freeß-hCG 0.41 LOW T18 (Edwards' syndrome) 1:100000 Low Risk HIGH PAPP-A 0.62 T13 (Patau syndrome) 1:34000 Low Risk LOW HIGH

# **INTERPRETATION**

The First Trimester Screening for the given sample is found SCREEN NEGATIVE.

# SUGGESTIONS AND OTHER FINDINGS

In view of free bHCG MoMs observed in the mother, focused serial survillance for assessment of fetal growth can be considered.



UK NEQAS

Lab Reg. No. 90968

Verified by

Verified by **Mr. Pradip Kadam** Incharge Biochemistry



Verified by **Dr. Suresh Bhanushali** MD (Path), Consultant Pathologist Page 1 of 3

Lilac Insights Pvt. Ltd. 301-302, Building A-1, Rupa Solitaire Millennium Business Park, MIDC Industrial Area, Sector-1, Navi Mumbai, Maharashtra 400710. Phone: +91 22 41841438; Website: www.lilacinsights.com; For queries or complaints, please email: info@lilacinsights.com | CIN - U85191MH2011PTC217513



Sample Type:Serum





Patient name : Mrs. BHAVANI RAMU

Sample ID: 2300170526

Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method:Elec	ctroche	emiluminescence								
				PREGNA	NCY DETAILS					
No. of fetuse	es	:1		EDD	:27/04/202	4 Ageat	<b>Term</b> : 26.9	Years		
GA is Based on		: CRL 63mm at 19/10/2023		LMP Date	:28/07/202	:28/07/2023 LMP Certainty : Reg		ular		
Smoking : None Parity :		Height	:	Weigh	Weight : 69.00 Kg					
Ethinicity:As	sian	FHR :								
Previous pregnancy history		Pre	Pre-eclampsia history		Other findings					
Down syndrome Edwards' syndrome			PE in	PE in previous pregnancy		Insulin dependent diabetes				
Patau syndrome					Pat. mother had PE		Chronic hypertension			
		Date   GA: Gestation Age		Menstrual Period	EHR: Fetal Heart Ra		<i>,</i> ,			
EDD. Estimate	u Duc L				f Birth			eciumpoia ( D O D. D ate		
SPECIMEN DETAILS										
Sample ID		:2300170526	CRL	: 63 mm	Test Name	Conc	. Unit	Corr. Mom		
Collection Date		: 19/10/2023	CRL2	:	Free-ß-hCC	<b>i</b> 13.23	3 ng/mL	0.41		
Scan Date		: 19/10/2023	BPD	:	NT	0.9	mm	0.64		
GA at Coll Date		: 12 Weeks 5 Days	BPD2	:	PAPP-A	2262.0	00 mIU/L	0.62		
GA at Scan Date		: 12 Weeks 5 Days	нс	:						
<b>Received on</b> : 23/10/2023 <b>HC2</b>		HC2	:							
GA: Gestation	n Age   C	RL: Crown Rump Lengtl	h   BPD: Bi-pa	arietal Diameter   I	HC: Head Circumfere	ence   free-ß-hCG: free	-Beta Human Cho	rionic Gonadotropin		
		NT:	Nuchal Trans	slucency   PAPP-A.	Pregnancy-associate	ed Plasma Protein-A				
				R	ISKS					
Disorder: Down Syndrome						Result:	Low Ris	k 🔵		
Final risk:	1:770	000	Age risk:	1:1300						
Cutoff	1:250	)	Risk type	e Risk At Ter	m					
Disorder: Edwards' Syndrome						Result:	Low Ris	k 🔵		
Final risk: 1:1000		0000	Age risk:	1:8400						
Cutoff 1:10		)	Risk type	e Risk At Ter	m					
Disorder: Patau Syndrome						Result:	Low Ris	k 🔵		
Final risk:	1:340		Age risk:	1:13000				-		
Cutoff	1:100	)	Risk type	Risk At Ter	·m					







Page 2 of 3

Lilac Insights Pvt. Ltd. 301-302, Building A-1, Rupa Solitaire Millennium Business Park, MIDC Industrial Area, Sector-1, Navi Mumbai, Maharashtra 400710. Phone: +91 22 41841438; Website: www.lilacinsights.com; For queries or complaints, please email: info@lilacinsights.com | CIN - U85191MH2011PTC217513







#### Patient name : Mrs. BHAVANI RAMU

# Sample ID: 2300170526

# PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

## PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

# High Risk

Low Risk

Intermediate

**High Risk or Screen Positive Result:** A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

**Intermediate Risk result:** An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

## SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

# For more information, visit our website at: <u>www.lilacinsights.com/faq-pns</u>

# DISCLAIMERS

#### Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
  further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
  abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT



Page 3 of 3