





To: Cancyte Technologies Pvt Ltd-Bangalore

1st Cross Road,

Shankarapuram Basavanagudi.

Karnataka

Bangalore - 560004

Contact:

Report Of: Mrs. SHREYA PAL

Pt. Contact: 9739074082

#### Sample ID 2310040057 Understand Your Report In Detail Patient ID 1102332966 Hosptial ID CANOBG230696 Received on 12/12/2023 18:33 Registered on 14/12/2023 11:16 Scan OR code Reported on Referred by Dr. Reena Dr. SAVITA SRIKANTH SHIRODKAR Sonography by

Patient DOB: 16/12/1994

## **EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT**

#### Patient Name: Mrs. SHREYA PAL

**EVIC** Screen\* is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probability of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

# UKNEQAS: United Kingdom National External Quality Assessment Service

RIQAS: Randox International Quality Assessment Scheme



The Risk Assessment Performed Using
CE-Marked Antenatal Risk Evaluation Software
Certified by the British Standards Institute
(BSI)- ISO 13485:2016

T21 (Down syndrome)	1:100000	Low Risk	LOW	INTERMEDIATE HIGH
T18 (Edwards' syndrome)	1: 100000	Low Risk	LOW	HIGH
T13 (Patau syndrome)	1: 100000	Low Risk	LOW	HIGH
Pre-eclampsia before 34 w	reeks <b>1:54</b>	High Risk	LOW	HIGH

# MULTIPLE OF MEDIAN (MoM)

Free ß-hCG	0.54	
AFP	1.27	
PAPP-A	2.29	
PLGF	1.03	

## **INTERPRETATION**

The First Trimester Enhanced Screening for the given sample is found SCREEN POSITIVE for Pre-eclampsia.









Page 1 of 3









Patient name: Mrs. SHREYA PAL Sample ID: 2310040057

**PREGNANCY DETAILS** 

Sample Type: Serum Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method: Time-resolved Fluroimmunoassay

No. of fetuses GA is Based on Smoking: None Ethinicity:Asian	:1 :CRL 68mm at 11/12/2023 Parity : Nulliparous FHR :	EDD LMP Date Height	: 17/06/2024 : 11/09/2023 : 160.0 cm	Age at Term LMP Certainty Weight	: 29.5 Years : Regular : 96.00 Kg	
-	pregnancy history	Pre-e	clampsia history	Oth	ner findings	
Down syndrome	Edwards' syndrome	PE in p	revious pregnancy	Insulin de	ependent diabetes	
Patau syndrome	NTD syndrome	Pat. mo	other had PE	Chronic	hypertension	

EDD: Estimated Due Date | GA: Gestation Age | LMP: Last Menstrual Period | FHR: Fetal Heart Rate | NTD: Neural Tube Defect | PE: Pre-eclampsia | DOB: Date of Birth

# SPECIMEN DETAILS

Sample ID	: 2310040057	CRL	: 68 mm	Test Name	Conc.	Unit	Corr. Mom
Collection Date	: 11/12/2023	CRL2	:	Free-ß-hCG	11.10	ng/mL	0.54
Scan Date	: 11/12/2023	BPD	:	NB	Present		
GA at Coll Date	: 13 Weeks 0 Days	BPD2	:	AFP	17.10	U/mL	1.27
GA at Scan Date	: 13 Weeks 0 Days	HC	:	NT	1.3	mm	0.87
Received on : 12/12/2023	HC2	:	PAPP-A	5910.00	mU/L	2.29	
			PLGF	61.86	pg/mL	1.03	
				MAP	113.33	mmHg	1.11
				UTPI	1.55		1.07

GA: Gestation Age | CRL: Crown Rump Length | BPD: Bi-parietal Diameter | HC: Head Circumference | free-ß-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency | PAPP-A: Pregnancy-associated Plasma Protein-A

			RISKS			
	own Syndrome			Result:	Low Risk 🛑	
Final risk:	1:100000	Age risk:	1:1019			
Cutoff	1:250	Risk type	Risk At Term			
Disorder: Ed	wards' Syndrome			Result:	Low Risk	
Final risk:	1:100000	Age risk:	1:9166			
Cutoff	1:100	Risk type	Risk At Term			
Disorder: Pa	tau Syndrome			Result:	Low Risk	
Final risk:	1:100000	Age risk:	1:27524			
Cutoff	1:100	Risk type	Risk At Term			
Disorder: PE	Disorder: PE < 34 weeks		Result:	High Risk		
Final risk:	1:54					
Cutoff	1: 100	Risk type	Risk at Term			















Patient name: Mrs. SHREYA PAL Sample ID: 2310040057

### PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

#### PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

**High Risk or Screen Positive Result:** A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk **Intermediate Risk result:** An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

### SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: <u>www.lilacinsights.com/faq-pns</u>

### **DISCLAIMERS**

#### Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

## Note:

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- PE risk stratification is done using a cut-off of 1:100 as per ASPRE study.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT

Page 3 of 3



