



To: Add-On Scans and Labs-Bangalore

No 5, MGR Complex,

Marathahalli -Sarjapur Rd, Above Pizza Hut,

Kaikondrahalli

Karnataka

Bangalore - 560035

Contact:

Report Of: Mrs. USHA KUMARI

Pt. Contact: 1000000000



Dr. ASHWINI R.S Sonography by

Patient DOB: 17/08/1995

Sample ID 2410005803 Understand Your Report In Detail Patient ID 1102341564 Hosptial ID 240460029 Received on 15/02/2024 17:04 Registered on 15/02/2024 17:33 Reported on Referred by Dr. CHAITRA GOWDA

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. USHA KUMARI

EVIC Screen* is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK) guidelines for First Trimester Screening to determine the probality of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

UKNEQAS: United Kingdom National External Quality Assessment Service

RIQAS: Randox International Quality Assessment **Scheme**



The Risk Assessment Performed Using **CE-Marked Antenatal Risk Evaluation Software** Certified by the British Standards Institute (BSI)- ISO 13485:2016

T21 (Down syndrome) 1: 910 Intermediate Risk LOW INTERMEDIATE HIGH T18 (Edwards' syndrome) 1: 100000 Low Risk LOW HIGH PAPP-A 0.93	RISK ASSESSMENT					MULTIPLE OF MEDIAN (MoM)
T18 (Edwards' syndrome) 1: 100000 Low Risk Low HIGH T13 (Patau syndrome) 1: 70000 Low Risk Low HIGH Pro-estampsia before 34 weeks 1:480	T21 (Down syndrome)	1:910	Intermediate Risk	LOW	INTERMEDIATE HIGH	
Pro eclampcia before 34 weeks 1:480	T18 (Edwards' syndrome)	1: 100000	Low Risk	LOW	HIGH	
Pre-eclampsia before 34 weeks 1:480 Low Risk Low HIGH	T13 (Patau syndrome)	1:70000	Low Risk	LOW	HIGH	
	Pre-eclampsia before 34 wee	ks 1:480	Low Risk	LOW	HIGH	

INTERPRETATION

The First Trimester Screening for the given sample is found Intermediate Risk for Down Syndrome.

SUGGESTIONS AND OTHER FINDINGS

- In view of intermediate risk (Risk between 1:251 to 1:1000), further counselling is recommended.
- Latest guidelines suggest further evaluation of intermediate risk patients by the following options as indicated:
- a) Integrated screening with detailed Genetic Sonogram (Detection rate: 92-95%), ref: Kypros Nicolaides et al, Fetal Diagn Ther 2014;35:174-184.

b) Non-Invasive Prenatal Testing/ Screening (NIPT) (Detection rate: >99%), ref: ISPD guidelines 2015.

c) Definitive testing through Fetal Karyotyping.

In view of free bHCG MoMs observed in the mother, kindly consider correlation with fetal growth and well being scan at 28 - 30 weeks.





Verified by Mr. Pradip Kadam Incharge Biochemistry (FMF ID: 147760)



MD (Path), Consultant Pathologist





Patient name: Mrs. USHA KUMARI Sample ID: 2410005803 Sample Type: Serum Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald Method: Electrochemiluminescence **PREGNANCY DETAILS** No. of fetuses **EDD** :27/08/2024 Age at Term : 29.0 Years :1 GA is Based on : CRL 57.7mm at 15/02/2024 **LMP Date** :23/11/2023 LMP Certainty: Regular Smoking: None Parity: Nulliparous Height : 158.0 cm Weight :60.20 Kg Ethinicity: Asian **FHR Previous pregnancy history** Pre-eclampsia history Other findings Down syndrome Edwards' syndrome PE in previous pregnancy Insulin dependent diabetes Patau syndrome NTD syndrome Pat. mother had PE Chronic hypertension EDD: Estimated Due Date | GA: Gestation Age | LMP: Last Menstrual Period | FHR: Fetal Heart Rate | NTD: Neural Tube Defect | PE: Pre-eclampsia | DOB: Date of Birth **SPECIMEN DETAILS** Sample ID :2410005803 **CRL** :57.7 mm **Test Name** Conc. Unit Corr. Mom Free-ß-hCG 306.60 8.88 **Collection Date** : 15/02/2024 CRL2 ng/mL NB Present Scan Date : 15/02/2024 **BPD** BPD2: NT 1.6 mm 1.23 **GA at Coll Date** : 12 Weeks 2 Days mIU/L PAPP-A 3390.00 0.93 **GA at Scan Date** HC : 12 Weeks 2 Days 0.92 MAP 76.70 mmHg Received on : 15/02/2024 HC2 UTPI 1.57 1.00 GA: Gestation Age | CRL: Crown Rump Length | BPD: Bi-parietal Diameter | HC: Head Circumference | free-\(\beta\)-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency | PAPP-A: Pregnancy-associated Plasma Protein-A **RISKS** Disorder: Down Syndrome Result: Intermediate Risk 1:910 Final risk: Age risk: 1:1100 Cutoff 1:250 Risk type Risk At Term Disorder: Edwards' Syndrome Result: Low Risk Final risk: 1:100000 Age risk: 1:7800 Cutoff 1:100 Risk type Risk At Term Disorder: Patau Syndrome Result: Low Risk



Final risk:

Final risk:

Cutoff

Cutoff

1:70000

1:100

1:480

1:100

Disorder: PE < 34 weeks



1:12000

Risk At Term

Risk at Term

Age risk:

Risk type

Risk type



Result:

Low Risk





Patient name: Mrs. USHA KUMARI Sample ID: 2410005803

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Low Risk

Low Risk or Screen Negative Result: A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: <u>www.lilacinsights.com/faq-pns</u>

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- PE risk stratification is done using a cut-off of 1:100 as per ASPRE study.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT



