



To: Dr.Veena Suresh-Thane shree darshan apartment ground floor Maharashtra - 400605 Contact: 9819137649 Report Of: Mrs. BANDANA SAHANI Pt. Contact: 8356953520		Sample ID Patient ID Received on Registered on Reported on Referred by Sonography by	2400200116 10024100466 15/11/2024 16:17 15/11/2024 16:24 - Dr. Veena Suresh Dr. Vaibhav Gaikwad	Understand Your Report In Detail
---	--	--	---	-------------------------------------

EVICOSCREEN - EVIDENCE BASED COMPREHENSIVE PRENATAL SCREENING REPORT

Patient Name: Mrs. BANDANA SAHANI

Patient DOB: 10/07/1997

EVIC Screen" is an evidence based prenatal screening program curated by Lilac Insights in accordance with the Fetal Medicine Foundation (UK)

guidelines for First Trimester Screening to determine the probality of most common chromosomal aneuploidies in a pregnancy. It utilizes:

- Hormonal values from the pregnancy measured on Fetal Medicine foundation (UK) accredited analyzers and reagents
- Robust indigenous medians from over 7 lac+ pregnancies for different gestation ages
- Risk calculations from evidence based algorithms validated through large international studies

UKNEQAS: United Kingdom National External Quality The Risk Assessment Performed Using Assessment Service **CE-Marked Antenatal Risk Evaluation Software Certified by the British Standards Institute RIQAS:** Randox International Quality Assessment (BSI)- ISO 13485:2016 Scheme **RISK ASSESSMENT 1ULTIPLE OF** MEDIAN (MoM) T21 (Down syndrome) 1:5400 Low Risk LOW INTERMEDIATE HIGH Freeß-hCG 6.23 Low Risk T18 (Edwards' syndrome) 1:100000 LOW HIGH PAPP-A 1.81 T13 (Patau syndrome) 1:65000 Low Risk LOW HIGH **INTERPRETATION** The First Trimester Screening for the given sample is found SCREEN NEGATIVE.

SUGGESTIONS AND OTHER FINDINGS

In view of free bHCG MoMs observed in the mother, kindly consider correlation with fetal growth and well being scan at 28 - 30 weeks.

JK NEQAS Lab Reg. No. 90968

Verified by Mr. Pradip Kadam Incharge Biochemistry (FMF ID: 147760)

Brede

Verified by **Dr. Suresh Bhanushali** MD (Path), Consultant Pathologist

Swetter -=

Page 1 of 3





Patient name : Mrs. BANDANA SAHANI

Sample ID : 2400200116

Sample	e Type:Serum
Sampro	

Risk assessment: Algorithm validated by SURUSS 2003, N.J Wald

Method:Chemilun	ninescence								
			PREGNANC	CY DETAILS					
No. of fetuses	:1		EDD	: 30/05/2025	Age at Terr	n :27.8	Years		
GA is Based on	: CRL 52mm at 14/	/11/2024	LMP Date	: 15/08/2024	LMP Certa	inty :Regu	ılar		
Smoking : None	Parity :		Height	:	Weight	:49.0	0 Kg		
Ethinicity:Asian	FHR :								
Previous pregnancy history		Pre-eclampsia history		Other findings					
Down syndro	me 🔲 Edwards's	syndrome	PE in pre	evious pregnancy	Insul	in depende	nt diabetes		
Patau syndrome 🔲 NTD syndrome		Pat. mother had PE		Chronic hypertension					
EDD: Estimated Due	Date GA: Gestation Age	/LMP: Last Me	enstrual Period FH	IR: Fetal Heart Rate NTD:	Neural Tube Defe	ect PE: Pre-e	clampsia DOB: Date		
			ofBi	irth					
SPECIMEN DETAILS									
Sample ID	:2400200116	CRL :	52 mm	Test Name	Conc.	Unit	Corr. Mom		
Collection Date	: 15/11/2024	CRL2 :		Free-ß-hCG	316.31	ng/mL	6.23		
Scan Date	: 14/11/2024	BPD :		NB	Present				
GA at Coll Date	: 12 Weeks 0 Days	BPD2 :		NT	1.5	mm	1.26		
GA at Scan Date	: 11 Weeks 6 Days	HC :		PAPP-A	9814.44	mIU/L	1.81		
Received on	: 15/11/2024	HC2 :							
GA: Gestation Age CRL: Crown Rump Length BPD: Bi-parietal Diameter HC: Head Circumference free-ß-hCG: free-Beta Human Chorionic Gonadotropin NT: Nuchal Translucency PAPP-A: Pregnancy-associated Plasma Protein-A									
			RIS	KS					
Disorder: Down Syndrome R		Res	ult:	Low Risl	<				
Final risk: 1:54	00	Age risk:	1:1200						
Cutoff 1:25	0	Risk type	Risk At Term						
Disorder: Edwards	' Syndrome			Res	sult:	Low Risl	< •		



Final risk:

Final risk:

Cutoff

Cutoff

1:100000

1:100

1:65000

1:100

Disorder: Patau Syndrome



Age risk:

Risk type

Age risk:

Risk type

Verified by Mr. Pradip Kadam Incharge Biochemistry (FMF ID: 147760)

1:8200

1:12000

Risk At Term

Risk At Term

Verified by **Dr. Suresh Bhanushali** MD (Path), Consultant Pathologis Low Risk

Result:





Sample ID: 2400200116

Patient name : Mrs. BANDANA SAHANI

PRENATAL SCREENING BACKGROUND

Every pregnant woman carries a certain degree of risk that her fetus/baby may have certain chromosomal defect/ abnormalities. Diagnosis of these fetal chromosomal abnormalities requires confirmatory testing through analysis of amniocytes or Chorionic Villous Samples (CVS). However, amniocentesis and CVS procedures carry some degree of risk for miscarriage or other pregnancy complications (Tabor and Alfirevic, 2010). Therefore in routine practice, prenatal screening tests are offered to a pregnant woman to provide her a personalised risk for the most common chromosomal abnormalities (T21-Down syndrome, T18- Edwards' syndrome, T13- Patau syndrome) using her peripheral blood sample. Based on this risk assessment, if the risk is high or intermediate, you can take informed decision of opting for invasive procedure such as amniocentesis or CVS followed by confirmatory diagnostic test(s), as per discussion with your clinician.

PRENATAL SCREENING TESTS ARE NOT CONFIRMATORY TESTS. THEY ARE LIKELIHOOD ASSESSMENT TESTS.

You may get your prenatal screening result as either of the following:-

High Risk

Low Risk

High Risk or Screen Positive Result: A High Risk Result does not mean that the pregnancy is affected with the condition. It means that the likelihood of the pregnancy having a condition is higher than the cut-off (Most commonly used cut-off is 1:250 and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis).

Later and the term

and this represents the risk of pregnancy loss from confirmatory testing through CVS or amniocentesis). **Low Risk or Screen Negative Result:** A Low Risk result does not mean that the pregnancy is not affected with a condition. It means that the likelihood of the pregnancy having a condition is lower than the cut-off.

Intermediate Risk result: An intermediate Risk result means that the pregnancy has an equivocal or a borderline risk of being affected with a condition. In this case, you may want to choose a second stage screening modality like an Integrated Screening Test that is done between 16 to 20 weeks of pregnancy or a Non-invasive Prenatal Screening Test between 12 to 20 weeks of pregnancy before taking a decision on an invasive confirmatory testing. This will help you improve the sensitivity of the screening test keeping an invasive test a last option were you to come as a high risk in the second stage screening test.

SIGNIFICANCE OF MULTIPLE OF MEDIANS (MoMs)

Prenatal Screening determines the likelihood of the pregnancy being affected with certain conditions by analysing levels of certain hormones. These hormones are Feto placental products (released by Fetus or placenta). Their levels not only indicate propensity of the fetus being affected with certain chromosomal conditions, they also provide indication of placental insufficiency that can potentially lead to pregnancy complications like Pre-Eclampsia or Intra-Uterine Growth Restriction. It is therefore important to take cognisance of the Reported MoMs alongside the Risk results.

For more information, visit our website at: <u>www.lilacinsights.com/faq-pns</u>

DISCLAIMERS

Limitations of the Test:

As prenatal screening tests are not confirmatory diagnostic tests, the possibility of false positive or false negative results can not be denied. The results issued for this test does not eliminate the possibility that this pregnancy may be associated with other chromosomal or sub- chromosomal abnormalities, birth defects and other complications.

Nuchal Translucency is the most prominent marker in screening for Trisomy 13, 18, 21 in the first trimester and should be measured in accordance with the Fetal Medicine Foundation (UK) guidelines. Nuchal Translucency or Crown Rump Length measurement, if not performed as per FMF (UK) imaging guidelines may lead to erroneous risk assessments and Lilac Insights bears no responsibility for errors arising due to sonography measurements not performed as per these criteria defined by international bodies such as FMF (UK), ISUOG.

It is assumed that the details provided along with the sample are correct. The manner in which this information is used to guide patient care is the responsibility of the healthcare provider, including advising for the need for genetic counselling or additional diagnostic testing like amniocentesis or Chorionic Villus Sampling. Any diagnostic test should be interpreted in the context of all available clinical findings. As with any medical test, there is always a chance of failure or error in sample analysis though extensive measures are taken to avoid these errors.

Note:

- Quality of the Down syndrome screening program (Biochemical values, MoMs and Risk assessments) is monitored by UKNEQAS on an ongoing basis.
- This interpretation assumes that patient and specimen details are accurate and correct.
- Lilac Insights does not bear responsibility for ultrasound measurements like CRL,NT,NB etc. We strongly recommend that ultrasound measurements are performed as per FMF (UK)/ISUOG practice guidelines.
- It must be clearly understood that the results represent risk and not diagnostic outcomes. Increased risk does not mean that the baby is affected and
 further tests must be performed before a firm diagnosis can be made. A Low Risk result does not exclude the possibility of Down's syndrome or other
 abnormalities, as the risk assessment does not detect all affected pregnancies.
- Each sample received at Lilac Insights' processing centre is handled with the utmost sensitivity and care. All samples received on Sundays and National holidays are stored as per specific guidelines for the respective specimens and processed on the next day.

END OF REPORT

